

USING ‘INVISIBLE’ DECAY MODES AS PROBES OF Z' COUPLINGS

J.L. HEWETT and T.G. RIZZO

*High Energy Physics Division, Argonne National Laboratory,
Argonne, IL 60439, USA*

ABSTRACT

We explore the possibility that Z' couplings can be probed using decay modes which involve neutrinos once Standard Model backgrounds are directly determined by the data itself. For some models, sufficient statistics are available at either the SSC or LHC to render these modes useful for coupling determinations, provided the mass of the Z' is not much larger than 1 TeV, if we assume other new physics background sources are absent.

If a new neutral gauge boson, Z' , is discovered at the SSC or LHC, an immediate goal will be the determination of its couplings to fermion pairs. Knowledge of these couplings will allow us to ascertain the extended electroweak model from which the Z' has originated. At an e^+e^- collider, such as LEP, the SLC, or the NLC, this is a relatively straightforward procedure that has already been successfully employed for the Standard Model (SM) Z . Performing a similar task in the hadron supercollider environment is far from easy and has prompted much activity in the theoretical and detector development arenas during the past three years¹. All of the techniques suggested in Ref. 1 are somewhat limited in scope so that new tools are continuously being sought to aid in the determination of Z' couplings at hadron supercolliders. In this report we outline some preliminary results from an analysis of Z' decay modes which involve neutrinos. As is always the case, the greatest difficulty in the use of missing E_t or p_t final states is the size of the SM backgrounds. In the two cases examined here, however, we may be able to employ the data itself to precisely determine these backgrounds. To be as specific as possible, we limit our discussion to only a few extended gauge models: (i) the Left-Right Symmetric Model (LRM)², (ii) the Alternative Left-Right Model (ALRM)³, (iii) a model where the Z' couples exactly like the SM Z (SSM), and (iv) the E_6 -inspired Effective Rank-5 Model (ER5M)⁴. In the latter example, the model contains an additional parameter, θ , which lies in the range $-90^\circ < \theta < 90^\circ$. Specific choices of this parameter correspond to special models discussed in the literature, *i.e.*, $\psi(\theta = 0^\circ)$, $\chi(\theta = -90^\circ)$, and $\eta(\theta = 37.76^\circ)$.

Our first analysis concerns the quantity $r_{\nu\nu Z} \equiv \Gamma(Z' \rightarrow Z\nu\bar{\nu})/\Gamma(Z' \rightarrow \ell^+\ell^-)$ which has been suggested by several of the authors of Ref. 1 as a probe of the Z' couplings. However, as shown explicitly in our earlier work, the SM background (B)

Model	SSC		LHC	
	2.5	0.5	2.5	0.5
SSM	2.794	0.664	0.656	0.213
LRM	0.344	0.083	0.079	0.026
ALRM	1.172	0.249	0.314	0.091
ψ	0.202	0.047	0.048	0.015
χ	0.838	0.214	0.178	0.063
η	0.090	0.020	0.022	0.007

Table 1. Values of the parameter f for the SSC and LHC assuming either value of the rapidity cut ($\eta \leq 2.5$ or 0.5) for several different extended electroweak models discussed in the text.

from ZZ production (with one of the Z 's decaying to $\nu\bar{\nu}$) can be up to several orders of magnitude larger than the $Z' \rightarrow Z\nu\bar{\nu}$ signal (S) in the absence of sophisticated cuts. This is demonstrated in Figs. 1a and 1b, where the missing p_t distributions are presented for both processes (signal and background) at the SSC and LHC, subject to mild rapidity (η) cuts, assuming a Z' mass of 1 TeV. For the various choices of extended models, the dotted curve in these figures must be scaled by a factor, f , which is model and rapidity cut dependent and is given numerically in Table 1. Clearly, unless the background can be precisely determined, there is very little hope of seeing the Z' signal even if both the η and p_t cuts are optimized. However, we observe that the SM process $pp \rightarrow ZZ \rightarrow Z\ell^+\ell^-$ can be precisely *measured* for a fixed p_t of the $\ell^+\ell^-$ pair, subject to the constraint that the pair invariant mass reconstructs to a Z . If this set of data were then to be rescaled by the ratio of the $Z \rightarrow \nu\bar{\nu}$ to $Z \rightarrow \ell^+\ell^-$ branching fractions, and corrected for acceptances, the background for the Z' process in question can then be determined. As a check of this procedure, the $Z\nu\bar{\nu}$ data normalization can be examined at low p_t where the Z' contribution is negligible. Since the background peaks at low p_t and the signal is absent at $p_t \gtrsim 500$ GeV for a 1 TeV Z' , demanding $p_t^{min} \leq p_t \leq 500$ GeV, with p_t^{min} in the 50-200 GeV range, will significantly enhance $S/\sqrt{B}$. We find that a value of $p_t^{min} = 200$ GeV gives the largest $S/\sqrt{B}$ for both choices of the rapidity cut.

To proceed further, we simulate data for both the signal and the background within the framework of a given model, subject to the cuts above, and determine the size of the excess sample of events obtained by integrating the corresponding missing p_t distributions. Since the size of the background has been anticipated from the charged lepton data as discussed above, it is then subtracted from the total number of events. If a statistically significant residual is obtained, it can be attributed to a 1 TeV Z' . (We remind the reader that the Z' will already have been discovered and its mass will have been determined via the charged lepton decay modes before the technique we're discussing here is applied.) Normalizing this excess to the number of $\ell^+\ell^-$ events in the Z' discovery channel eliminates the systematic uncertainties due to, *e.g.*, structure function variations and the integrated luminosity determination. Table 2 displays the inverse of the relative error in $r_{\nu\nu Z}$, *i.e.*, $(\Delta r_{\nu\nu Z}/r_{\nu\nu Z})^{-1}$ (which is equivalent to $S/\sqrt{B}$), obtained from this method for a number of extended gauge models. Here, the resulting

Model	SSC		LHC	
	2.5	0.5	2.5	0.5
SSM	8.379	4.104	10.70	6.704
LRM	1.032	0.520	1.289	0.818
ALRM	3.515	1.561	5.123	2.864
ψ	0.608	0.295	0.783	0.472
χ	2.518	1.342	2.904	1.983
η	0.270	0.125	0.359	0.220

Table 2. Values of the quantity $(\Delta r_{\nu\nu Z}/r_{\nu\nu Z})^{-1}$ at the SSC and LHC assuming either choice of the rapidity cut.

statistical and systematic errors have been combined in quadrature. The integrated luminosity ($\mathcal{L}$) for the SSC(LHC) was assumed to be 10(100) fb^{-1} in obtaining these relative errors which have magnitudes that are found to scale approximately as $1/\sqrt{\mathcal{L}}$. From the Table we see that at least for some of the extended models, the value of $r_{\nu\nu Z}$ can be sufficiently well determined as to provide some additional information about the nature of the Z' couplings. It is clear from this analysis, however, that this technique will most likely fail for larger Z' masses due to the loss in statistics necessary to observe the signal relative to the background.

A second approach to obtaining information about the Z' coupling to $\nu\bar{\nu}$ is to examine the production of a high p_t jet in association with both the Z' as well as the SM Z . The essential idea is an extension of the monojet analysis by the CDF Collaboration⁵ from which they obtained a constraint on the number of neutrinos. We consider the following four processes: $pp \rightarrow Z + jet, Z' + jet \rightarrow \ell^+\ell^- + jet, \nu\bar{\nu} + jet$ at both the SSC and LHC as functions of the jet transverse momentum, assuming $|\eta_j| < 0.5$ or 2.5. (In practice, the η_j cut is chosen to be 2.5 since the lower value only results in decreased statistics.) The two processes which involve charged leptons in the final state can be directly measured, with extremely high statistics in the case of the SM Z , and are easily separable by demanding the dilepton pairs lie in invariant mass bins associated with the Z or the Z' . (We again assume that the Z' mass is already well determined from the discovery channel; for purposes of demonstration we take its value to be 1 TeV.) From the number of observed SM Z induced $\ell^+\ell^-j$ events we can deduce the anticipated number of SM Z monojet events, which we treat as a background, through again simply scaling by the well measured ratio of $Z \rightarrow \nu\bar{\nu}$ to $Z \rightarrow \ell^+\ell^-$ branching fractions and correcting for acceptance differences. Of course, the monojet rate also obtains a contribution from the $Z' + jet$ intermediate state, which is proportional to the $Z' \rightarrow \nu\bar{\nu}$ branching fraction, and may appear as a signal over the SM Z background. For a given extended electroweak model, we simulate both the signal and the backgrounds for a minimum jet p_t in the 50 – 200 GeV range allowing for statistical fluctuations in the number of events. We subtract from the total event sample, after integrating over η and p_t , the anticipated number of SM events, obtained through the rescaling technique mentioned above and normalize any potential excess by the number of Z' induced $\ell^+\ell^-j$ events. If we assume that other new physics sources

of monojets are not present, this normalized excess of events is directly proportional to the quantity $R_\nu = \Gamma(Z' \rightarrow \nu\bar{\nu})/\Gamma(Z' \rightarrow \ell^+\ell^-)$, which is quite sensitive to Z' coupling variations. (Assuming that invisible decays of the Z' arise solely from its decay into the 3 generations of SM left-handed ν 's.) In taking the ratio above we again have eliminated potentially large systematic errors arising from variations in the parton distributions and integrated luminosity uncertainties.

What range of values can we expect for the ratio R_ν ? In extended models where the new generator which couples to the Z' commutes with the SM $SU(2)_L$ generators, we find that R_ν can be expressed as $R_\nu = 3c_L^2/(c_L^2 + c_R^2)$, with $c_{L(R)}$ being the left(right)-handed coupling of the charged lepton to the Z' . Clearly, in this case, we must have $0 \leq R_\nu \leq 3$, independently of the specific choice of model. When $R_\nu = 0(3)$, the $Z'\ell^+\ell^-$ coupling is purely right(left)-handed. This ‘commuting’ condition is readily satisfied by the LRM, ALRM, and ER5M choices but *not* for the SSM case since the $SU(2)_L$ generators do not commute amongst themselves. Figs. 2a-c show the anticipated values of R_ν in the ER5M case, and the associated errors which follow from this analysis, as functions of θ at both the SSC and LHC with various cuts on the jet transverse momentum. We see immediately that (i) R_ν is quite sensitive to the value of θ (thus demonstrating its usefulness as a model discriminator), (ii) a stricter p_t cut reduces the size of the anticipated errors, and (iii) the same cuts applied to the data from the SSC will yield smaller errors than that similarly obtained at the LHC even when the factor of 10 difference in integrated luminosity is accounted for. In Fig. 3a, we see the corresponding results for the LRM as a function of $\kappa \equiv g_R/g_L$, the ratio of the $SU(2)_R$ to $SU(2)_L$ gauge couplings. For large values of κ we note that the couplings of the charged leptons become purely right-handed. As in the case of the parameter θ in the ER5M, R_ν is quite sensitive to the value of κ , particularly when it lies in its natural range, *i.e.*, $0.55 \leq \kappa \leq 1$, anticipated by GUT analyses⁶. Fig. 3b displays a comparison of the values of R_ν obtained for the ALRM, SSM, LRM with $\kappa = 1$, and ER5M as a function of θ . It is clear from the figure that R_ν can be determined with reasonable precision and may be an extremely useful discriminator of extended gauge models provided the assumptions we have made above are valid. The presence of monojets from other new physics sources, such as SUSY, will render this approach extremely difficult if not impossible. As in the previous analysis we observe that for substantially larger Z' masses there will be insufficient statistics to overcome the large SM backgrounds.

In the analyses above we have considered two possible ways of gaining information on the couplings of a Z' to neutrinos in the face of large SM backgrounds. Rather than relying solely on theoretical calculations, which suffer from many uncertainties, we propose instead to use data from SSC and/or LHC experiments themselves to determine the size of these backgrounds. In following this route and always employing ratios of numbers of events, many of the systematic uncertainties are found to cancel. For the first possibility examined, the case of three-body Z' decays, we found that it may be possible to resurrect the quantity $r_{\nu\nu Z}$ as an model discriminator provided strong p_t cuts and invariant mass binning can be applied to the data without appreciable loss in rate. For many extended electroweak models, this technique was found to work reasonably well for a Z' in the 1 TeV range, but for significantly larger masses, there is a dramatic

loss in statistical power. A second technique that employs potential excesses in monojet events as a signal for $Z' \rightarrow \nu\bar{\nu}$, is extremely dependent upon several assumptions, in particular, that no other non-SM sources of monojets, such as SUSY, exist. If such sources are absent (or possibly well determined by other means), it was found that the ratio of the partial widths of the Z' to $\nu\bar{\nu}$ and $\ell^+\ell^-$ can be reasonably well determined for Z' masses in the 1 TeV range.

Acknowledgements

This work has been supported by the U.S. Department of Energy, Division of High Energy Physics under contract W-31-109-ENG-38.

References

1. A. Fiandrino and P. Taxil, *Phys. Rev.* **D44**, 3409 (1991) and *Phys. Lett.* **B293**, 242 (1992); J.D. Anderson, M.H. Austern, and R.N. Cahn, *Phys. Rev. Lett.* **69**, 25 (1992) and *Phys. Rev.* **D46**, 290 (1992); F. del Aguila et al., *Phys. Rev.* **D48**, 425 (1993); J.L. Hewett and T.G. Rizzo, *Phys. Rev.* **D45**, 161 (1992), **D47**, 4981 (1993), and Argonne National Laboratory report ANL-HEP-PR-92-33 (1992); T.G. Rizzo, *Phys. Rev.* **D47**, 956 (1993); see also, T.G. Rizzo, Argonne National Laboratory report ANL-HEP-PR-93-18 (1993); M. Cvetič and P. Langacker, *Phys. Rev.* **D46**, 14 (1992) and **D46**, 4943 (1992); M. Cvetič, P. Langacker, and B. Kayser, *Phys. Rev. Lett.* **68**, 2871 (1992); ATLAS Collaboration Letter of Intent, D. Gingrich *et al.*, CERN Report LHC/I2, October, 1992; A. Henriques and L. Poggioli, ATLAS Collaboration Note PHYS-NO-010, October, 1992.
2. For a review of the LRM and original references, see R.N. Mohapatra, *Unification and Supersymmetry*, (Springer, New York, 1986); P. Langacker and S. Uma Sankar, *Phys. Rev.* **D40**, 1569 (1989).
3. E. Ma, *Phys. Rev.* **D36**, 274 (1987) and *Mod. Phys. Lett.* **A3**, 319 (1988); K.S. Babu *et al.*, *Phys. Rev.* **D36**, 878 (1987); V. Barger and K. Whisnant, *Int. J. Mod. Phys.* **A3**, 879 (1988); J.F. Gunion *et al.*, *Int. J. Mod. Phys.* **A2**, 118 (1987); T.G. Rizzo, *Phys. Lett.* **B206**, 133 (1988).
4. For a review of such models, see J.L. Hewett and T.G. Rizzo, *Phys. Rep.* **183**, 193 (1989).
5. R. Markeloff, CDF Collaboration, talk given at the *Division of Particles and Fields Meeting*, Fermilab, Nov. 1992.
6. See, for example, N.G. Deshpande, E. Keith, and T.G. Rizzo, *Phys. Rev. Lett.* **70**, 3185 (1993).

Fig. 1: Missing p_t distribution for the background process, $pp \rightarrow 2Z \rightarrow Z\nu\bar{\nu}$, at the SSC(solid) and the LHC(dash-dotted) as well as for the signal(dotted), $pp \rightarrow Z' \rightarrow Z\nu\bar{\nu}$, assuming an η cut of (a)2.5 or (b)0.5.

Fig. 2: The ratio R_ν as a function of the parameter θ in the ER5M assuming a Z' mass of 1 TeV with $|\eta_j| < 2.5$ at (a)the SSC with $p_t > 200$ GeV, or (b) with $p_t > 50$ GeV. In either case, the errors reflect the combined statistical and systematic errors associated with an integrated luminosity of $10fb^{-1}$. (c) Same as (a) but for the LHC with an integrated luminosity of $100fb^{-1}$.

Fig. 3: (a)Same as Fig. 2a but for the LRM as a function of the parameter $\kappa \equiv g_R/g_L$. (b) A comparison of the values for R_ν as in Fig. 2a for the ALRM, SSM, LRM with $\kappa = 1$, and ER5M as a function of θ .

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9309207v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9309207v1>

This figure "fig1-3.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9309207v1>